



## SPECIFICATION

DRUGS FOR MEDICAL USE ENABLING NUCLEAR  
MAGNETIC RESONANCE DIAGNOSIS BY SCALAR COUPLINGRECEIVED  
AUG 23 2002  
TECHCENTER 160012900

10

## TECHNICAL FIELD

The present invention relates to drugs for medical use which can be detected by the nuclear magnetic resonance method, particularly to drugs for medical use which contain a compound that exerts a relaxation effect on the proton of  $^1\text{H}$  hydrogen atom in system by scalar coupling.

## BACKGROUND ART

It is acknowledged by everybody that medicines confer great benefits to mankind. However, medicines are not essential components for the maintenance of life and, in the case of a medicine which is a foreign material to a living body, it is not provided with a mechanism of being selectively transported to a target organ or tissue in the body unlike essential components for a living body. Consequently, the medicine administered into a living body comes to distribute itself not only to the intended site but to everywhere in the living body.

Accordingly, many medicines are not only inefficient in efficacy, but cause side effects in greater or lesser extent. Moreover, since the disease state is not uniform for each patient, the distribution state of the medicine becomes

more complicated, and it is difficult, for each patient, to select a truly effective medicine, to predict its effect and to estimate the prognosis including ~~side effect~~ side effects.

For example, generally the "normal dose" in therapeutic

5 practice is based on the results of clinical trials conducted for only several hundred cases or the results accumulated by the intuition and experience of doctors. Thus, at the present state, there can be found no objective means for suppressing unfavorable effects resulting from unnecessary distribution of  
10 a medicine and bringing out the intended effective action thereof to the greatest extent for each patient. That is to say, the problem is that it is yet unknown how and to what extent a medicine is transported to a target organ or tissue.

Medical treatment of the present day is supported by  
15 many peripheral sciences and technologies. Particularly marked is the progress of imaging diagnosis including X-ray diagnosis, nuclear magnetic resonance diagnosis, ultrasonic diagnosis and nuclear medicine diagnosis. Furthermore, various drugs for diagnostic use suited to each of the diagnostic  
20 methods are being developed, and it has become possible to determine externally the in vivo pharmacokinetics, such as circulation and distribution of these drugs for diagnostic use in real time. In such a case, however, though previous drugs for diagnostic use enable accurate specification of the morbid  
25 site in the living body and identification of the morbid state based on the in vivo pharmacokinetics thereof, they give virtually no suggestion, after the diagnosis, for the proper selection of therapeutic drugs to be actually used and for the

prediction of their efficacy. Nevertheless, in recent years, for example, as a contrast medium for positron emission tomography (PET), one of the nuclear medicine diagnoses, 2-fluoro( $^{18}\text{F}$ )-2-deoxy-D-glucose, which is a glucose-analogue, appears at the actual spots of medical treatment. However, though it is structurally analogous to glucose, which is an existing drug, it does not reflect the in vivo pharmacokinetics of glucose in a strict sense; further, the object of its use is limited to the diagnosis of tumors or the like, and it is not used as the means for collecting information by using glucose as a drug. Moreover, it has the disadvantage of being radioactive.

Among the several imaging diagnostic methods which enable non-invasive observation of the inside of a living body from the outside, the diagnostic method with nuclear magnetic resonance has a number of merits which are not observed in other methods of imaging diagnosis. In particular, since it gives a high contrast between soft tissues as compared with previous X-ray CT, it has a very high discriminating ability between various soft tissues such as the ones in the brain, heart, liver, kidney and the like. Furthermore, it permits tomography in any desired direction and can provide blood flow information; these points are favorable in tracing in vivo behavior of drugs. Moreover, the apparatus for nuclear magnetic resonance has already been used in many medical organizations, so that the method is not restricted in its use unlike PET, for which the apparatus is provided only to a

limited number of medical organizations.

The principle of the diagnostic method with nuclear magnetic resonance is as follows: when a high frequency pulse including the resonance frequency of the hydrogen atom  $^1\text{H}$  is  
5 irradiated, a resonance phenomenon takes place at the hydrogen nuclei and the resonance signal is observed. This serves for imaging the state of distribution of the proton of hydrogen present as water in the living body. It is also apparent that the method is free from the risk of radiation exposure unlike  
10 the X-ray diagnostic method and nuclear medicine diagnostic method.

It has been reported that an element having a quadruple nucleus of a nuclear spin of 1 or more, for example  $^{17}\text{O}$ , shortens the transverse relaxation time ( $T_2$ ), which forms  
15 the basis in imaging protons, by scalar coupling with the proton of hydrogen atom  $^1\text{H}$  bonded thereto (S. Meiboom, J. Chem. Phys., 39, 375, 1961). Arai et al. have disclosed a method wherein, for the purpose of imaging the distribution of water  $\text{H}_2^{17}\text{O}$  formed as a part of the metabolite of  $^{17}\text{O}_2$  by using the  
20 above-mentioned technology,  $^{17}\text{O}_2$  is mixed with a perfluoro compound and an emulsifier and administered into a living body (Japanese National Publication (Kohyo) 3-500896). However, Arai et al., have mentioned as the result of their later study that the change of signal intensity due to  $^{17}\text{O}_2$  metabolism and  
25 the condition of metabolic function do not always correspond well to each other, and a good imaging of  $\text{H}_2^{17}\text{O}$  of the metabolite cannot be attained (JP-A-6-22936)

Further, some attempts have been made to obtain the

image of a proton in high sensitivity by utilizing scalar coupling. For example, Navon et al. have developed a measuring method using  $\text{H}_2^{17}\text{O}$  with  $^{17}\text{O}$  irradiation (US5479924).

## 5 DISCLOSURE OF THE INVENTION

In order to overcome the above-mentioned problems that the medical science on individual difference which forms the basis for the differentiation of patients mainly in the therapeutical aspect has not yet been able to establish  
10 scientifically, an object of the present invention is to provide a physiologically acceptable medical drug which enables external detection of the effective circulation or distribution of the drug, used for a disease, in the target organ or tissue in vivo where it is needed by the nuclear  
15 magnetic resonance method before or at the same time as the administration of a therapeutic agent to each patient. Said medical drug can be used in the nuclear magnetic resonance imaging, the nuclear magnetic resonance spectrometry or the determination of relaxation time.

20 The present inventors have made extensive study to attain the above-mentioned object. As a result, the inventors have found that a medical drug can itself become a pharmacokinetic-diagnostic agent which can provide for each patient information on the circulation and distribution of the  
25 drug in vivo by a nuclear magnetic resonance method utilizing scalar coupling when the drug contains, among the constituents contained in the drug, a compound which carries in its chemical structure at least one member selected from the group consisting of -OH, -NH and -SH.

Thus, the present invention provides a drug for medical use characterized by containing a compound which carries in its chemical structure at least one member selected from the group consisting of  $^{-17}\text{OH}$ ,  $^{14}\text{NH}$  and  $^{-33}\text{SH}$ , wherein the  
5 above  $^{17}\text{O}$ ,  $^{14}\text{N}$  or  $^{33}\text{S}$  exerts a relaxation effect on the proton of hydrogen  $^1\text{H}$  bonded thereto and the relaxation effect spreads through the exchange of a proton in a vital component of a target organ or tissue in a living body with the above-mentioned proton, thus enabling detection by the nuclear  
10 magnetic resonance method.

The above-mentioned vital component of a target organ or tissue in a living body is usually water, but it may also be lactic acid, N-acetylaspartic acid, etc.

The "detection by the nuclear magnetic resonance  
15 method" allows for measurement of phenomenon wherein the  $^{17}\text{O}$ ,  $^{14}\text{N}$  or  $^{33}\text{S}$  in the above compound contained in the above-mentioned medical drug exerts a relaxation effect on the proton of hydrogen  $^1\text{H}$  bonded thereto and then the proton exchanges itself with a proton in a vital component of a  
20 target organ or tissue in a living body by means of nuclear magnetic resonance imaging, nuclear magnetic resonance spectrometry or relaxation time measurement each using a proton as the detection nucleus.

According to the present invention, it becomes  
25 possible to provide a medical drug which enables external detection of the effective circulation or distribution of the drug, used for a disease, in the target organ or tissue in vivo where it is needed by the nuclear magnetic resonance

method before the administration of a therapeutic agent to each patient or in real time, and it becomes further possible to establish the medical science on individual difference which forms the basis for differentiation mainly in the  
5 therapeutic aspect.

#### MODE FOR CARRYING OUT THE INVENTION

The drug for medical use according to the present invention is selected from drugs which contain as a  
10 constituent a compound having in its chemical structure at least one member selected from the group consisting of the -OH, -NH and -SH groups. Furthermore, the whole or a part of the O, N or S atoms constituting the respective groups are substituted with their respective isotopes  $^{17}\text{O}$ ,  $^{14}\text{N}$  or  $^{33}\text{S}$ .  
15 Accordingly, said drug for medical use can be synthesized by using a raw material or intermediate, by which the -OH, -NH or -SH group is to be introduced, in which the whole or a part of the O, N or S atoms have been substituted with their respective stable isotopes  $^{17}\text{O}$ ,  $^{14}\text{N}$  or  $^{33}\text{S}$ , according to a known  
20 method for preparing a medical drug. Though  $^{17}\text{O}$  exists in nature only in a low concentration of 0.04%, the separation of  $^{17}\text{O}$  and the enrichment of  $^{16}\text{O}$  by  $^{17}\text{O}$  themselves are not the object of the present invention. With regard thereto, several methods are described in literature which include, for example,  
25 fractional distillation of heavy water, electrolysis and laser (isotope) separation.

The selection of the medical drug of the present invention containing a compound, in which the whole or a part

of the O, N or S atoms of the -OH, -NH or -SH groups are substituted with their respective stable isotopes  $^{17}\text{O}$   $^{14}\text{N}$  or  $^{33}\text{S}$  can be made by those skilled in the art within the range not deleterious to the object of the present invention, and any  
5 desired existing medical drug can be used according to the therapeutical or diagnosis object. More specifically, selection may be made as desired according to the disease state of each patient from therapeutic agents, such as drugs for the central nervous system, drugs for the circulatory  
10 system, drugs for the digestive system, drugs for the urogenital system and drugs for tumors; nutritional or tonic agents; agents for blood and humor such as infusion; or agents for diagnosis, such as X-ray contrast media, MRI contrast media, ultrasonic contrast media and radiopharmaceuticals.  
15 Since  $^{17}\text{O}$  and  $^{33}\text{S}$ , which are each a stable isotope element, have respectively exactly the same chemical properties as those of usual oxygen  $^{16}\text{O}$  and sulfur  $^{32}\text{S}$ ,  $^{17}\text{O}$  and  $^{33}\text{S}$  do not show pharmacokinetics of a different nature also in a living body. The above-mentioned compound to be substituted may be any of  
20 the active ingredients of medical drugs, additives and solvents. Specifically, the active ingredient is preferably a sugar, particularly glucose, amino acid, etc; the solvent is preferably an aqueous solvent, particularly water. They may be compounded according to the formulation ratio of respective  
25 medical drugs. The dosage form may also be selected according to the respective medical drugs and may be either a solution or a lyophilized product which is dissolved before use. Further, it may have been processed with a material for a drug



delivery system, such as a liposome or the like.

The medical drug thus obtained is administered according to the administration routes determined for respective medical drugs. It is administered, for example, intravenously, intraarterially, intramuscularly or orally, but it may be administered precutaneously as the occasion demands. When it is desired to select a proper medical drug or to estimate its efficacy, the medical drug of the present invention may be administered prior to the full-scale administration of a medical drug. When it is desired, for example, to monitor the administration of a therapeutic agent and whether a medical drug is properly circulated or distributed in a target organ or tissue, simultaneously, the medical drug of the present invention may be used as the whole or a part of the medical drug.

The dose of the medical drug of the present invention may be appropriately selected according to the use of the drug, the degree of enrichment of  $^{17}\text{O}$ ,  $^{14}\text{N}$  or  $^{33}\text{S}$ , and the kind of the nuclear magnetic resonance method used as the means of determination. The nuclear magnetic resonance method used as the means of determination may be any desired one so long as it is a nuclear magnetic resonance method using a proton as the detection nucleus. Nuclear magnetic resonance imaging, nuclear magnetic resonance spectrometry or relaxation time determination is preferable. Nuclear magnetic resonance imaging is particularly preferred. In an experiment to determine the distribution of  $\text{H}_2^{17}\text{O}$  ( $^{17}\text{O}$  content: about 89%) using a rat cerebral ischemic model conducted by the present

inventors as an experimental example , an image based on tissue infusion difference was obtained by the T2-weighted spin echo method (nuclear magnetic resonance imaging using a proton as the detection nucleus), which is a common imaging method in medical diagnosis. Further, for example, nuclear magnetic resonance imaging using  $^{17}\text{O}$  irradiation is more preferably used since a more enhanced sensitivity can be obtained.

For example, in the percutaneous local therapy (PEIT) of hepatic cancer, wherein pure ethanol is injected percutaneously and transhepatically into hepatic cancer cells to coagulate and necrotize the cancer cells, there is a high possibility of the ethanol diffusing excessively to cause coagulation and necroses even of the normal hepatic cells. In such a case, for example, by injecting the present medical drug obtained by substituting the whole or a part of the  $^{16}\text{OH}$  group of ethanol with  $^{17}\text{O}$ , under a nuclear magnetic resonance apparatus, and observing the resulting image, it becomes possible to provide a therapy which limits the injection range to only cancer cells. Sodium hyaluronate, which is administered into cavitas articulare in the case of osteoarthritis of the knee, can be used, for example, for similar purposes.

On the other hand, in the case of an infusion, which is used as the substitute of a blood transfusion from the viewpoint of replenishing water, it is important how it spreads throughout a living body. In the case of an electrolyte infusion, which can be of many varieties according

to the kind and the concentration of the electrolyte contained therein, a proper drug must be selected for each patient to improve the disease state. In this instance, by using the present drug in which the whole or a part of water as a solvent has been substituted with  $^{17}\text{O}$ , it becomes possible to judge an electrolyte infusion and determine what composition is to be used for improving the disease state of each patient.

#### EXAMPLES

The present invention is described in detail below with reference to the Examples, but the technical scope of the present invention is not limited thereby.

##### Example 1 Synthesis of $[3\text{-}^{17}\text{OH}]\text{glucose}$

20 g (0.07 mol) of methyl-4,6-O-benzylidene- $\alpha$ -D-allopyranoside was dissolved in 108 ml of pyridine to obtain a pale yellow solution. While the solution was being cooled, 43 g (0.23 mol) of p-toluenesulfonyl chloride was added thereto over 15 minutes or more, and the resulting mixture was stirred at  $30^{\circ}\text{C}$  for 48 hours. The reaction solution turned light brown and formed a white precipitate. The reaction solution was poured into ice water and extracted with chloroform. The chloroform layer was separated and washed successively with 5% sulfuric acid, 4% aqueous sodium hydrogen carbonate solution and water. Then the chloroform layer was dried over  $\text{MgSO}_4$  and evaporated to dryness. The yellow sirupy residue was recrystallized from ethanol to obtain 32 g of a white solid. Yield 79%.

31 g (0.05 mol) of the 2,3-bis(O-p-toluenesulfonyl)

derivative obtained above was dissolved in 350 ml of chloroform, and a mixture of 35 ml of a 28% sodium methoxide methanol solution and 65 ml of methanol was added thereto. The resulting mixture was kept at room temperature for 48 hours while being gently stirred and then diluted with 300 ml of water. The chloroform layer was separated, washed twice with water, then dried over  $\text{MgSO}_4$  and evaporated to dryness. The residue obtained was crystallized from chloroform-ether to obtain a white solid.

10           The 500 ml of THF was added to 5.29 g (20 mmols) of methyl-2, 3-anhydro-4, 6-O-benzylidene- $\alpha$ -allopyranoside obtained above. While the resulting mixture was being stirred, 1 g of Nafion-H and 1 g of water ( $^{17}\text{O}$  content: 10%) were slowly added thereto, and the mixture was stirred overnight at room temperature. The insolubles and Nafion-H were removed by suction filtration and the filtrate was evaporated to dryness under reduced pressure to obtain a white solid. The white solid was hydrolyzed to obtain 1.4 g of  $[3-^{17}\text{OH}]$  glucose.

20   Example 2 Preparation of agent for humor stoperative restoring liquid) using water ( $^{17}\text{O}$ ) as solvent

          The titled drug was prepared according to the formulation of "KN replenisher 4A" (a trade name, mfd. by Otsuka Pharmaceutical Factory, Inc.), which is a postoperative restoring liquid used as the replenishing agent for water and electrolytes for the whole body early after the operation. Water ( $^{16}\text{O}$ ) was added to 0.234 g of sodium chloride, 8.002 g of

glucose and sodium lactate to make up the total to 100 ml. To a 0.5 ml portion thereof was added an equal quantity of water containing 10.5% of  $^{17}\text{O}$  so as to prepare an  $^{17}\text{O}$  water-containing KN replenisher 4A ( $^{17}\text{O}$  content: 5.25%)

5

### Example 3 Phantom imaging of aqueous [1- $^{17}\text{OH}$ ] glucose solution

Water was added to [1- $^{17}\text{OH}$ ] glucose available on the market so as to prepare a solution of a concentration of 62.55  
10 mg/ml. As a control, glucose ( $^{16}\text{O}$ ) was used to prepare a solution of a concentration of 62.9 mg/ml. These solution solutions were respectively sealed into glass tubes and imaged with a nuclear magnetic resonance imaging apparatus of 2 tesla, Omega CSI. When the echo time was set at 200 msec, the  
15 signal intensity ratios of the aqueous [1- $^{17}\text{OH}$ ] glucose solution and the aqueous ( $^{16}\text{O}$ ) glucose solution were respectively 79.3 and 93.1. Thus, it has become apparent that [1- $^{17}\text{OH}$ ] glucose can be a compound which exerts a relaxation effect by scalar coupling on the water protons of the tissue  
20 in which they are distributed.

### Example 4 Determination of relaxation time of body liquid drug (postoperative restoring liquid) using water ( $^{17}\text{O}$ ) as solvent

25 The relaxation time of the ( $^{17}\text{O}$ ) water- containing KN replenisher 4A ( $^{17}\text{O}$  content: 5.25%) obtained in Example 2 was determined. As the control, a KN replenisher 4A ( $^{17}\text{O}$  content: natural abundance ratio) prepared by using water ( $^{16}\text{O}$ ) alone as

the solvent was used. The relaxation time was determined by the CPMG method using a Model JNM-FSE-60 pulse NMR. As the result, the relaxation time of the  $^{17}\text{O}$  water-containing KN replenisher 4A was  $181.19 \pm 0.50$  msec, being significantly shortened as compared with the relaxation time,  $626.36 \pm 1.01$  msec, of the  $^{16}\text{O}$  water-containing replenisher 4A. From the result, it has become apparent that an effective medical drug  $^{17}\text{O}$  water-containing KN replenisher 4A which could exert a relaxation effect on a water proton by scalar coupling could be prepared.

Example 5 Determination of relaxation time of pure ethanol ( $^{17}\text{OH}$ ) in water

A 0.2 ml portion of pure ethanol ( $^{17}\text{OH}$ ,  $^{17}\text{O}$  content: 10%) available on the market was mixed with 0.2 ml of water ( $^{16}\text{O}$ ) (50% in terms of ethanol concentration). As a control, a solution was prepared by mixing pure ethanol ( $^{16}\text{OH}$ ) in the same manner as above. The relaxation times of the two samples thus prepared were determined by the CPMG method using a Model JNM-FSE-60 pulse NMR. As the result, the relaxation time of the pure ethanol ( $^{17}\text{OH}$ ) dilution water was  $420.22 \pm 0.99$  msec, being significantly shortened as compared with the relaxation time ( $779.12 \pm 4.91$  msec) of pure ethanol ( $^{16}\text{OH}$ ) of the control. From the result, it is apparent that pure ethanol ( $^{17}\text{OH}$ ) can become

a medical drug which, after administration and distribution, exerts a relaxation effect on the water proton of a tissue by scalar coupling.